

References

Lemon Balm

- 3
- Tyler VE. Herbs of Choice. Binghamton, NY: Pharmaceutical Products Press, 1994.
- 285
- Bisler H, Pfeifer R, Kluken N, Pauschinger P. [Effects of horse-chestnut seed on transcapillary filtration in chronic venous insufficiency]. Dtsch Med Wochenschr 1986;111:1321-9. [View abstract](#)
- 790
- Wolbling RH, Leonhardt K. Local therapy of herpes simplex with dried extract from *Melissa officinalis*. Phytomedicine 1994;1:25-31.
- 4912
- Electronic Code of Federal Regulations. Title 21. Part 182 -- Substances Generally Recognized As Safe. Available at:
<https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/CFRSearch.cfm?CFRPart=182>
- 7049
- Holtmann G, Madisch A, Juergen H, et al. A double-blind, randomized, placebo-controlled trial on the effects of an herbal preparation in patients with functional dyspepsia [Abstract]. Ann Mtg Digestive Disease Week 1999 May.
- 9993
- Akhondzadeh S, Noroozian M, Mohammadi M, et al. *Melissa officinalis* extract in the treatment of patients with mild to moderate Alzheimer's disease: a double blind, randomised, placebo controlled trial. J Neurol Neurosurg Psychiatry 2003;74:863-6. [View abstract](#).
- 9994
- Kennedy DO, Scholey AB, Tildesley NT, et al. Modulation of mood and cognitive performance following acute administration of *Melissa officinalis* (lemon balm). Pharmacol Biochem Behav 2002;72:953-64. [View abstract](#).
- 9995
- Koytchev R, Alken RG, Dundarov S. Balm mint extract (Lo-701) for topical treatment of recurring herpes labialis. Phytomedicine 1999;6:225-30. [View abstract](#).
- 10421
- St-Onge MP, Jones PJ. Physiological effects of medium-chain triglycerides: potential agents in the prevention of obesity. J Nutr 2002;132:329-32.. [View abstract](#).
- 10422
- Burns A, Byrne J, Ballard C, Holmes C. Sensory stimulation in dementia. BMJ 2002;325:1312-3.. [View abstract](#).
- 10423
- Cerny A, Shmid K. Tolerability and efficacy of valeren/lemon balm in healthy volunteers (a double blind, placebo-controlled, multicentre study). Fitoterapia 1999;70:221-8.
- 12724
- Madisch A, Holtmann G, Mayr G, et al. Treatment of functional dyspepsia with a herbal preparation. A double-blind, randomized, placebo-controlled, multicenter trial. Digestion 2004;69:45-52. [View abstract](#).
- 13089
- Melzer J, Rosch W, Reichling J, et al. Meta-analysis: phytotherapy of functional dyspepsia with the herbal drug preparation STW 5 (Iberogast). Aliment Pharmacol Ther 2004;20:1279-87. [View abstract](#).
- 14416
- Muller SF, Klement S. A combination of valeren and lemon balm is effective in the treatment of restlessness and dyssomnia in children. Phytomedicine 2006;13:383-7. [View abstract](#).
- 14618
- Lee J, Chae K, Ha J, et al. Regulation of obesity and lipid disorders by herbal extracts from *Morus alba*, *Melissa officinalis*, and *Artemisia capillaris* in high-fat diet-induced obese mice. J Ethnopharmacol 2008;115:263-70. [View abstract](#).
- 16735

- Savino F, Cresi F, Castagno E, et al. A randomized double-blind placebo-controlled trial of a standardized extract of Matricariae recutita, Foeniculum vulgare and Melissa officinalis (ColiMil) in the treatment of breastfed colicky infants. *Phytother Res* 2005;19:335-40. [View abstract](#). **19405**
- Kennedy DO, Little W, Haskell CF, et al. Anxiolytic effects of a combination of *Melissa officinalis* and *Valeriana officinalis* during laboratory induced stress. *Phytother Res*. 2006 Feb;20:96-102. [View abstract](#). **19417**
- Lindahl O, Lindwall L. Double blind study of a valeren preparation. *Pharmacol Biochem Behav*. 1989 Apr;32:1065-6. [View abstract](#). **19427**
- Albrecht M, Berger W, Laux P, Schmidt U, et al. Psychopharmaka und Verkehrssicherheit. Der Einfluß von Euvegal® - Dragees forte auf die Fahrtüchtigkeit und Kombinationswirkungen mit Alkohol Z Allg Med 1995;71:1215-25. **19521**
- Vogt M, Tausch I, Wölbling RH, and et al. [Melissa extract in herpes simplex: a double-blind placebo-controlled study]. *Der Allgemeinarzt* 1991;13:832-841. **19522**
- Burns A, Perry E, Holmes C, et al. A double-blind placebo-controlled randomized trial of *Melissa officinalis* oil and donepezil for the treatment of agitation in Alzheimer's disease. *Dement Geriatr Cogn Disord*. 2011;31:158-64. [View abstract](#). **19523**
- Ballard CG, O'Brien JT, Reichelt K, Perry EK. Aromatherapy as a safe and effective treatment for the management of agitation in severe dementia: the results of a double-blind, placebo-controlled trial with *Melissa*. *J Clin Psychiatry* 2002 Jul;63:553-8. [View abstract](#). **19524**
- Kennedy DO, Little W, Scholey AB. Attenuation of laboratory-induced stress in humans after acute administration of *Melissa officinalis* (Lemon Balm). *Psychosom Med*. 2004 Jul-Aug;66:607-13. [View abstract](#). **19525**
- PARDO-ALDAVE K, DIAZ-PIZAN ME, VILLEGAS LF, et al. Child behaviour modulation during first dental visit after administration of lemon balm, Poster Sessions. *International Journal of Paediatric Dentistry* 2009;19(1):66-170. **19526**
- Büchner KH, Hellings H, Huber M, et al. [Double blind study as evidence of the therapeutic effect of Melissengeist on psycho-vegetative syndromes (author's transl)]. *Med Klin*. 1974 Jun 7;69:1032-6. [View abstract](#). **19528**
- Schmidt, U, Krieger, W, Frerick, H, and Schenk, N. Psychosomatische und psychische Störungen / Baldrian und Melisse statt synthetischer Psychopharmaka. 1992;14:15-19. **19529**
- Cases J. Leaf extract in the treatment of volunteers suffering from mild-to-moderate anxiety disorders and sleep disturbances. *Mediterr J Nutr Metab*. 2010;4(3):211-218. **19530**
- Kennedy DO, Wake G, Savelev S, et al. Modulation of mood and cognitive performance following acute administration of single doses of *Melissa officinalis* (Lemon balm) with human CNS nicotinic and muscarinic receptor-binding properties. *Neuropsychopharmacology*. 2003 Oct;28:1871-81. [View abstract](#). **19531**
- Chakurski I, Matev M, Koichev A, et al. [Treatment of chronic colitis with an herbal combination of *Taraxacum officinale*, *Hipericum perforatum*, *Melissa officinaliss*, *Calendula officinalis* and *Foeniculum vulgare*]. *Vutr Boles*. 1981;20:51-4. [View abstract](#). **19532**
- Madisch A, Melderis H, Mayr G, et al. [A plant extract and its modified preparation in functional dyspepsia. Results of a double-blind placebo controlled comparative study]. *Z Gastroenterol* 2001;39(7):511-7. [View abstract](#). **19534**

- Solberg E. The effects of powdered fertilized eggs on depression. *J Med Food*. 2011 Jul-Aug;14:870-5. [View abstract](#). **19535**
- Vejdani R, Shalmani HR, Mir-Fattah M, et al. The efficacy of an herbal medicine, Carmint, on the relief of abdominal pain and bloating in patients with irritable bowel syndrome: a pilot study. *Dig Dis Sci*. 2006 Aug;51:1501-7. [View abstract](#). **19536**
- Dressing H, Kohler S, and Muller WE. Improvement in sleep quality with a high dose valeren-melissa preparation . *Psychopharmakotherapie* 1996;3:123-130. **19537**
- Widy-Tyszkiewicz, E and Schminda, R. A randomized double blind study of sedative effects of phytotherapeutic containing valerian, hops, balm and motherwort versus placebo. *Herb Polon* 1997;2:154-159. **19538**
- Melzer J, Schrader E, Brattström A, et al. Fixed herbal drug combination with and without butterbur (Ze 185) for the treatment of patients with somatoform disorders: randomized, placebo-controlled pharmaco-clinical trial. *Phytother Res*. 2009 Sep;23:1303-8. [View abstract](#). **19701**
- Weizman Z, Alkrinawi S, Goldfarb D, et al. Efficacy of herbal tea preparation in infantile colic. *J Pediatr* 1993;122(4):650-652. [View abstract](#). **19723**
- Herberg, KW. Nebenwirkungen pflanzlicher Beruhigungsmittel/ Leistung und Befinden nach Einnahme einer Baldrian-Hopfen-Kombination. *Z.Allg Med* 1996;72:234-240. **19724**
- Chung MJ, Cho SY, Bhuiyan MJ, et al. Anti-diabetic effects of lemon balm (*Melissa officinalis*) essential oil on glucose- and lipid-regulating enzymes in type 2 diabetic mice. *Br.J.Nutr.* 2010;104:180-188. [View abstract](#). **19725**
- Soulimani R, Fleurentin J, Mortier F, et al. Neurotropic action of the hydroalcoholic extract of *Melissa officinalis* in the mouse. *Planta Med*. 1991 Apr;57:105-9. [View abstract](#). **19726**
- Sourgens H, Winterhoff H, Gumbinger HG, et al. Antihormonal effects of plant extracts. TSH- and prolactin-suppressing properties of *Lithospermum officinale* and other plants. *Planta Med*. 1982 Jun;45:78-86. [View abstract](#). **19727**
- Auf'mkolk M, Ingbar JC, Amir SM, et al. Inhibition by certain plant extracts of the binding and adenylate cyclase stimulatory effect of bovine thyrotropin in human thyroid membranes. *Endocrinology*. 1984 Aug;115:527-34. [View abstract](#). **19728**
- Santini F, Vitti P, Ceccarini G, et al. In vitro assay of thyroid disruptors affecting TSH-stimulated adenylate cyclase activity. *J Endocrinol Invest*. 2003 Oct;26:950-5. [View abstract](#). **30020**
- Meolie, A. L., Rosen, C., Kristo, D., Kohrman, M., Gooneratne, N., Aguillard, R. N., Fayle, R., Troell, R., Townsend, D., Claman, D., Hoban, T., and Mahowald, M. Oral nonprescription treatment for insomnia: an evaluation of products with limited evidence. *J Clin.Sleep Med* 4-15-2005;1(2):173-187. [View abstract](#). **49322**
- Wong, A. H., Smith, M., and Boon, H. S. Herbal remedies in psychiatric practice. *Arch Gen.Psychiatry* 1998;55(11):1033-1044. [View abstract](#). **58045**
- Hohmann, J., Zupko, I., Redei, D., Csanyi, M., Falkay, G., Mathe, I., and Janicsak, G. Protective effects of the aerial parts of *Salvia officinalis*, *Melissa Officinalis* and *Lavandula angustifolia* and their constituents against enzyme-dependent and enzyme-independent lipid peroxidation. *Planta Med* 1999;65(6):576-578. [View abstract](#). **59322**

Perry, E. K., Pickering, A. T., Wang, W. W., Houghton, P. J., and Perry, N. S. Medicinal plants and Alzheimer's disease: from ethnobotany to phytotherapy. *J Pharm Pharmacol* 1999;51(5):527-534. [View abstract](#).

59324

Wake, G., Court, J., Pickering, A., Lewis, R., Wilkins, R., and Perry, E. CNS acetylcholine receptor activity in European medicinal plants traditionally used to improve failing memory. *J Ethnopharmacol* 2000;69(2):105-114. [View abstract](#).

59326

Mikus, J., Harkenthal, M., Steverding, D., and Reichling, J. In vitro effect of essential oils and isolated mono- and sesquiterpenes on *Leishmania major* and *Trypanosoma brucei*. *Planta Med* 2000;66(4):366-368. [View abstract](#).

59327

Triantaphyllou, K., Blekas, G., and Boskou, D. Antioxidative properties of water extracts obtained from herbs of the species Lamiaceae. *Int J Food Sci Nutr* 2001;52(4):313-317. [View abstract](#).

59329

Mrljanova, M., Tekel'ova, D., Felklova, M., Reinoohl, V., and Toth, J. The influence of the harvest cut height on the quality of the herbal drugs *Melissae folium* and *Melissae herba*. *Planta Med* 2002;68(2):178-180. [View abstract](#).

59331

Gyllenhaal, C., Merritt, S. L., Peterson, S. D., Block, K. I., and Gochenour, T. Efficacy and safety of herbal stimulants and sedatives in sleep disorders. *Sleep Med Rev*. 2000;4(3):229-251. [View abstract](#).

59333

Araujo, C., Sousa, M. J., Ferreira, M. F., and Leao, C. Activity of essential oils from Mediterranean Lamiaceae species against food spoilage yeasts. *J Food Prot*. 2003;66(4):625-632. [View abstract](#).

59337

Patora, J., Majda, T., Gora, J., and Klimek, B. Variability in the content and composition of essential oil from lemon balm (*Melissa officinalis* L.) cultivated in Poland. *Acta Pol.Pharm.* 2003;60(5):395-400. [View abstract](#).

59338

Drozd, J. and Anuszewska, E. The effect of the *Melissa officinalis* extract on immune response in mice. *Acta Pol.Pharm.* 2003;60(6):467-470. [View abstract](#).

59339

Mimica-Dukic, N., Bozin, B., Sokovic, M., and Simin, N. Antimicrobial and antioxidant activities of *Melissa officinalis* L. (Lamiaceae) essential oil. *J Agric.Food Chem.* 5-5-2004;52(9):2485-2489. [View abstract](#).

59340

de Sousa, A. C., Alviano, D. S., Blank, A. F., Alves, P. B., Alviano, C. S., and Gattass, C. R. *Melissa officinalis* L. essential oil: antitumoral and antioxidant activities. *J Pharm.Pharmacol.* 2004;56(5):677-681. [View abstract](#).

59341

Gazola, R., Machado, D., Ruggiero, C., Singi, G., and Macedo, Alexandre M. *Lippia alba*, *Melissa officinalis* and *Cymbopogon citratus*: effects of the aqueous extracts on the isolated hearts of rats. *Pharmacol.Res* 2004;50(5):477-480. [View abstract](#).

59343

Ivanova, D., Gerova, D., Chervenkov, T., and Yankova, T. Polyphenols and antioxidant capacity of Bulgarian medicinal plants. *J Ethnopharmacol* 1-4-2005;96(1-2):145-150. [View abstract](#).

59344

Allahverdiyev, A., Duran, N., Ozguven, M., and Koltas, S. Antiviral activity of the volatile oils of *Melissa officinalis* L. against Herpes simplex virus type-2. *Phytomedicine*. 2004;11(7-8):657-661. [View abstract](#).

59346

Gaby, A. R. Natural remedies for Herpes simplex. *Altern.Med Rev*. 2006;11(2):93-101. [View abstract](#).

59350

- Awad, R., Levac, D., Cybulski, P., Merali, Z., Trudeau, V. L., and Arnason, J. T. Effects of traditionally used anxiolytic botanicals on enzymes of the gamma-aminobutyric acid (GABA) system. *Can.J Physiol Pharmacol.* 2007;85(9):933-942. [View abstract](#). **59351**
- Abuhamda, S., Huang, L., Elliott, M. S., Howes, M. J., Ballard, C., Holmes, C., Burns, A., Perry, E. K., Francis, P. T., Lees, G., and Chazot, P. L. Pharmacological profile of an essential oil derived from *Melissa officinalis* with anti-agitation properties: focus on ligand-gated channels. *J Pharm Pharmacol.* 2008;60(3):377-384. [View abstract](#). **59352**
- Soh, N. L. and Walter, G. Complementary medicine for psychiatric disorders in children and adolescents. *Curr.Opin.Psychiatry* 2008;21(4):350-355. [View abstract](#). **59355**
- Awad, R., Muhammad, A., Durst, T., Trudeau, V. L., and Arnason, J. T. Bioassay-guided fractionation of lemon balm (*Melissa officinalis* L.) using an in vitro measure of GABA transaminase activity. *Phytother.Res.* 2009;23(8):1075-1081. [View abstract](#). **59357**
- Raines, T., Jones, P., Moe, N., Duncan, R., McCall, S., and Ceremuga, T. E. Investigation of the anxiolytic effects of luteolin, a lemon balm flavonoid in the male Sprague-Dawley rat. *AANA.J.* 2009;77(1):33-36. [View abstract](#). **59358**
- Hanganu, D., Vlase, L., Filip, L., Sand, C., Mirel, S., and Indrei, L. L. The study of some polyphenolic compounds from *Melissa officinalis* L. (Lamiaceae). *Rev.Med.Chir Soc.Med.Nat.Iasi* 2008;112(2):525-529. [View abstract](#). **59359**
- Guginski, G., Luiz, A. P., Silva, M. D., Massaro, M., Martins, D. F., Chaves, J., Mattos, R. W., Silveira, D., Ferreira, V. M., Calixto, J. B., and Santos, A. R. Mechanisms involved in the antinociception caused by ethanolic extract obtained from the leaves of *Melissa officinalis* (lemon balm) in mice. *Pharmacol.Biochem.Behav.* 2009;93(1):10-16. [View abstract](#). **59360**
- Mencherini, T., Picerno, P., Russo, P., Meloni, M., and Aquino, R. Composition of the fresh leaves and stems of *Melissa officinalis* and evaluation of skin irritation in a reconstituted human epidermis model. *J.Nat.Prod.* 2009;72(8):1512-1515. [View abstract](#). **59362**
- Ibarra, A., Feuillere, N., Roller, M., Lesburgere, E., and Beracochea, D. Effects of chronic administration of *Melissa officinalis* L. extract on anxiety-like reactivity and on circadian and exploratory activities in mice. *Phytomedicine.* 2010;17(6):397-403. [View abstract](#). **59363**
- Kim, M. Y., Park, B. Y., Lee, H. S., Park, E. K., Hahm, J. C., Lee, J., Hong, Y., Choi, S., Park, D., Lee, H., and Yoon, M. The anti-angiogenic herbal composition Ob-X inhibits adipose tissue growth in obese mice. *Int.J.Obes.(Lond)* 2010;34(5):820-830. [View abstract](#). **59364**
- Taguchi, Y., Takizawa, T., Ishibashi, H., Sagawa, T., Arai, R., Inoue, S., Yamaguchi, H., and Abe, S. Therapeutic effects on murine oral candidiasis by oral administration of cassia (*Cinnamomum cassia*) preparation. *Nihon Ishinkin.Gakkai Zasshi* 2010;51(1):13-21. [View abstract](#). **59370**
- Ntalli, N. G., Ferrari, F., Giannakou, I., and Menkissoglu-Spirodi, U. Phytochemistry and nematicidal activity of the essential oils from 8 Greek Lamiaceae aromatic plants and 13 terpene components. *J.Agric.Food Chem.* 7-14-2010;58(13):7856-7863. [View abstract](#). **59377**
- Lara, M. S., Gutierrez, J. I., Timon, M., and Andres, A. I. Evaluation of two natural extracts (*Rosmarinus officinalis* L. and *Melissa officinalis* L.) as antioxidants in cooked pork patties packed in MAP. *Meat.Sci.* 2011;88(3):481-488. [View abstract](#). **59382**
- Yoon, M. and Kim, M. Y. The anti-angiogenic herbal composition Ob-X from *Morus alba*, *Melissa officinalis*, and *Artemisia capillaris* regulates obesity in genetically obese ob/ob mice. *Pharm.Biol.* 2011;49(6):614-619. [View abstract](#).

59384

Budzynska, A., Wieckowska-Szakiel, M., Sadowska, B., Kalembska, D., and Rozalska, B. Antibiofilm activity of selected plant essential oils and their major components. *Pol.J.Microbiol.* 2011;60(1):35-41. [View abstract](#).

59387

Perry, E. and Howes, M. J. Medicinal plants and dementia therapy: herbal hopes for brain aging? *CNS.Neurosci.Ther* 2011;17(6):683-698. [View abstract](#).

59390

Englberger, W., Hadding, U., Etschenberg, E., Graf, E., Leyck, S., Winkelmann, J., and Parnham, M. J. Rosmarinic acid: a new inhibitor of complement C3-convertase with anti-inflammatory activity. *Int J Immunopharmacol.* 1988;10(6):729-737. [View abstract](#).

59393

Kucera, L. S., Cohen, R. A., and Herrmann, E. C., Jr. Antiviral activities of extracts of the lemon balm plant. *Ann N.Y.Acad Sci* 7-30-1965;130(1):474-482. [View abstract](#).

59394

Kucera, L. S. and Herrmann, E. C., Jr. Antiviral substances in plants of the mint family (labiateae). I. Tannin of *Melissa officinalis*. *Proc Soc Exp Biol Med* 1967;124(3):865-869. [View abstract](#).

59395

Herrmann, E. C., Jr. and Kucera, L. S. Antiviral substances in plants of the mint family (labiateae). II. Nontannin polyphenol of *Melissa officinalis*. *Proc Soc Exp Biol Med* 1967;124(3):869-874. [View abstract](#).

59402

Larrondo, J. V., Agut, M., and Calvo-Torras, M. A. Antimicrobial activity of essences from labiates. *Microbios* 1995;82(332):171-172. [View abstract](#).

59404

Dimitrova, Z., Dimov, B., Manolova, N., Pancheva, S., Ilieva, D., and Shishkov, S. Antiherpes effect of *Melissa officinalis* L. extracts. *Acta Microbiol Bulg.* 1993;29:65-72. [View abstract](#).

59406

Yamasaki, K., Nakano, M., Kawahata, T., Mori, H., Otake, T., Ueba, N., Oishi, I., Inami, R., Yamane, M., Nakamura, M., Murata, H., and Nakanishi, T. Anti-HIV-1 activity of herbs in Labiateae. *Biol.Pharm Bull* 1998;21(8):829-833. [View abstract](#).

59407

Perry, E. K., Pickering, A. T., Wang, W. W., Houghton, P., and Perry, N. S. Medicinal plants and Alzheimer's disease: Integrating ethnobotanical and contemporary scientific evidence. *J Altern Complement Med* 1998;4(4):419-428. [View abstract](#).

59408

PARDO-ALDAVE, K. DIAZ-PIZAN M. E. VILLEGRAS L. F. BERNABE E. Child behaviour modulation during first dental visit after administration of lemon balm, Poster Sessions. *International Journal of Paediatric Dentistry* 2009;19(1):66-170.

59410

Holtmann, G., Madisch, A., and Juergen, H. A double-blind, randomized, placebo-controlled trial on the effects of an herbal preparation in patients with functional dyspepsia [Abstract]. *Ann Mtg Digestive Disease Week* 1999;May:A65.

59412

Hussain, A. I., Farooq Anwar, Nigam, P. S., Sarker, S. D., Moore, J. E., Rao, J. R., and Mazumdar, A. Antibacterial activity of some Lamiaceae essential oils using resazurin as an indicator of cell growth. *LWT - Food Science and Technology* 2012;44(4):1199-1206.

59417

Meftahizade, H., Sargsyan, E., and Moradkhani, H. Investigation of antioxidant capacity of *Melissa officinalis* L. essential oils. *Journal of Medicinal Plants Research* 2010;4(14):1391-1395.

59418

Rojas, J., Solís, H., and Palacios, O. In vitro anti-Trypanosoma cruzi activity of essential oils of ten medicinal plants. / Evaluación in vitro de la actividad anti Trypanosoma cruzi de aceites esenciales de diez plantas medicinales.1. *Anales de la Facultad de Medicina* 2010;71(3):161-165.

59420

Wolbling RH and Milbradt R. [Clinical manifestations and treatment of Herpes simplex infections]. Therapiewoche 1984;34:1193-1200.

59421

Wolbling RH and Leonhardt K. Local therapy of herpes simplex with dried extract from Melissa officinalis. Phytomedicine 1994;1:25-31.

59423

Dkbas, N., Bagc, E., Kotan, R., Cakmakc, R., Ozer, H., Mete, E., and Erdogan, G. Comparative antibacterial activities and chemical composition of some plants' oils against *Salmonella enteritidis*. Research on Crops 2010;11(1):118-124.

59425

Stanojevic, D., Ēomic', L., Stefanovic', O., and Sukdolak, S. S. In vitro synergistic antibacterial activity of *Melissa officinalis* L. and some preservatives. Spanish Journal of Agricultural Research 2010;8(1):109-115.

59426

Karimi, I., Hayatgheybi, H., Razmjo, M., Yousefi, M., Dadyan, A., and Hadipour, M. Anti-hyperlipidaemic effects of an essential oil of *Melissa officinalis* L. in cholesterol-fed rabbits. Journal of Applied Biological Sciences 2010;4(1):17-22.

59427

Cerny AS and Schmid K. Tolerability and efficacy of valerian/lemon balm in healthy volunteers; a double-blind placebo-controlled, multicentre study. Fitoterapia 1999;70(3):221-228.

59428

Dressing H. Valerian combination therapy vs. benzodiazepine: same efficacy in the treatment of sleeping disorders? Therapiewoche 1992;42(12):726-736.

59430

Buchner KH, Hellings H, Huber M, and et al. [Double blind study as evidence of the therapeutic effect of Melissengeist on psycho-vegetative syndromes]. Medizinische Klinik 1974;69(23):1032-1036.

59431

Schmidt, U, Krieger, W, Frerick, H, and Schenk, N. Psychosomatische und psychische Störungen / Baldrian und Melisse statt synthetischer Psychopharmaka. 1992;14:15-19.

59432

Lagoni, N. Wirksamkeitsprüfung eines pflanzlichen Tagessedativums in einer multizentrischen Studie. 1998;39(3):166-169.

59433

Dressing H, Kohler S, and Muller WE. Improvement in sleep quality with a high dose valerian-melissa preparation . Psychopharmakotherapie 1996;3:123-130.

59434

Widy-Tyszkiewicz, E and Schminda, R. A randomized double blind study of sedative effects of phytotherapeutic containing valerian, hops, balm and motherwort versus placebo. Herb Polon 1997;2:154-159.

59435

Orth-Wagner, S, Ressin, WJ, and Friedrich, I. Phytosedativum gegen Schlafstörungen / Klinische Wirksamkeit und Verträglichkeit eines Phytosedativums mit Auszügen aus Baldrianwurzel, Hopfenzapfen und Melissenblättern. 1995;16(147):156.

59437

Herberg, KW. Nebenwirkungen pflanzlicher Beruhigungsmittel/ Leistung und Befinden nach Einnahme einer Baldrian-Hopfen-Kombination. Z.Allg Med 1996;72:234-240.

59438

Cases J. Leaf extract in the treatment of volunteers suffering from mild-to-moderate anxiety disorders and sleep disturbances. Mediterr J Nutr Metab. 2010;4(3):211-218.

59441

Babpour, E., Angaji, S. A., and Angaji, S. M. Antimicrobial effects of four medicinal plants on dental plaque. Journal of Medicinal Plants Research 2009;3(3):132-137.

59442

Niculae, M., Spînu, M., Stefan, R., Sandru, C. D., Mircean, V., Brudasca, G. F., Bolfa, P., and Uricariu, R. In vitro efficacy of gentamicin and *Mellisa officinalis* essential oil against *Pseudomonas aeruginosa* isolated from dogs with otitis externa. Scientific Works - University of

Agronomical Sciences and Veterinary Medicine, Bucharest Series C, Veterinary Medicine
2010;56(3/4):280-284.

59446

Kizaibek, M., Kopp, B., Prinz, S., Popescu, R., and Upur, H. Antiproliferative activity of individual herbs of Abnormal Savda Munziq on HL-60 cells. Science & Technology Review 2009;19:94-98.

59447

Cohen RA, Kucera LS, and Herrmann EC. Antiviral activity of *Melissa officinalis* (lemon balm) extract (29600). Proceedings of the Society for Experimental Biology and Medicine 1964;117:431-434.

59460

Tittel G, Wagner H, and Bos R. [Chemical composition of the essential oil from *Melissa*]. Planta Medica 1982;46:91-98.

59478

Keskn, D., Oskay, D., and Oskay, M. Antimicrobial activity of selected plant spices marketed in the West Anatolia. International Journal of Agriculture and Biology 2010;12(6):916-920.

59481

Vogt M, Tausch I, Wölbling RH, and et al. [Melissa extract in herpes simplex: a double-blind placebo-controlled study]. Der Allgemeinarzt 1991;13:832-841.

59482

Borho, B. Biologische Therapie von funktionellen Magenerkrankungen. 1991;6:501-509.

91732

Alijaniha F, et al. Heart palpitation relief with *Melissa officinalis* leaf extract: double blind, randomized, placebo controlled trial of efficacy and safety. J Ethnopharmacol. 2015;164:378-384. doi: 10.1016/j.jep.2015.02.007. Epub 2015 Feb 11. [View abstract](#).

91733

Scholey A, et al. Anti-stress effects of lemon balm-containing foods. Nutrients. 2014;6(11):4805-4821. doi: 10.3390/nu6114805. [View abstract](#).

94122

Thompson A, Meah D, Ahmed N, et al. Comparison of the antibacterial activity of essential oils and extracts of medicinal and culinary herbs to investigate potential new treatments for irritable bowel syndrome. BMC Complement Altern Med. 2013;13:338. [View abstract](#).